arXiv:1212.6587v2 [nucl-th] 17 Feb 2013

# QGP formation time and direct photons from heavy ion collisions

Fu-Ming Liu[1, *]
[1]*Key laboratory of Quark and Lepton Physics (MoE) and Institute of Particle Physics,*
*Central China Normal University, Wuhan 430079, China*

(Dated: February 19, 2013)

The recent direct photon data, especially the large elliptic flow from AuAu collisions at $\sqrt{s_{NN}}$ = 200 GeV and PbPb collisions at $\sqrt{s_{NN}}$ =2.76 TeV has been investigated with hadronic data constrained 3+1D ideal hydrodynamical models. It turns out that those data support an earlier starting of the collective motion ($\tau_0 < 0.6$ fm/c) and a delayed formation of three-flavour QGP ( $\tau_{\rm QGP} \sim 2$ fm/c) in heavy ion collisions. The observed large elliptic flow of direct photon is still underestimated. A more realistic model with viscosity, an earlier $\tau_0$ and event-by-event fluctuation is expected to improve the work.

PACS numbers:

## I. INTRODUCTION

The recent direct photon data [1] from Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV shows a big enhancement at low transverse momentum ($p_{\rm t}$) region, similar to the PHENIX observation[2]. Also recently, a large elliptic flow ($v_2$) of direct photons, as large as that of hadrons, has been observed, both by PHENIX[3] and preliminarily by ALICE [4]. The large elliptic flow of direct photons challenges our understanding of heavy ion collisions.

To explain the large elliptic flow, some people consider additional sources of photon emission. For example, conformed anomaly as a source of soft photons [5] can produce a sizable elliptic flow. However, this addition source does not vanish at high $p_{\rm t}$ region and is ruled out by the PHENIX observation that $R_{AA} = 1$ at high $p_{\rm t}$ [6]. The contribution of pre-equilibrium Glasma stage to photon emission is estimated in [7], where a number of approximation must be made due to the current incomplete knowledge of this stage, which claimed this source may generate elliptic flow but hard to calculate. In [8], initial fluctuation has been considered and many useful qualitative results have been obtained.

Now we would like to check how well photon data and hadronic data can be simultaneously explained. Therefore the 3+1D ideal hydrodynamical model [9] is chosen for AuAu collisions at $\sqrt{s_{NN}} = 200$ GeV and [10] for PbPb collisions at $\sqrt{s_{NN}}$ =2.76 TeV. In both cases, hydro parameters are well fixed to reproduce excellently the bulk hadrons such as rapidity distribution, $p_{\rm t}$ spectra and elliptic flow. More extensive test on charm production and jet quenching, HBT and so on has been performed with those hydro models. The difference between the two models, *i.e.*, initial time $\tau_0 = 0.6$ fm/c for AuAu and $\tau_0 = 0.35$ fm/c for PbPb will tell us more than a single model chosen. Since data on bulk hadrons have been successfully explained, in this work we will focus on how well photon data can be explained.

Additionally, we would benefit from photon data and ask a simple question, "When is QGP formed in heavy ion collisions?" Here the QGP refers to a 3-flavour (locally) thermalized partons system to meet the observation of the saturated strangeness suppression factor $\gamma_s$ and free baryon number at midrapidity for central collisions [11].

Conventionally, this QGP matter is assumed to be formed at the beginning of collective motion, *i.e.*, the electromagnetic radiation from QGP are often calculated from the hydro initial time $\tau_0$ [12–14]. However, in heavy ion collisions at very low energies, no QGP formation but hydrodynamical model is still widely employed. In Landau hydrodynamical model [15], the hydro initial time is as early as just after the little bang, certainly QGP formation should be later since it takes time for the system to realize local equilibrium, both thermally and chemically. Hydro model is so widely employed, because the conervation law and the general relation $\epsilon = \epsilon(p)$ to close hydro problem, hold for many kinds of matter. But we should distinguish QGP formation time $\tau_{\rm QGP}$ from hydro initial time $\tau_0$.

How to estimate QGP formation time $\tau_{\rm QGP}$? The well-known thermal photon emission rates count photons emitted due to the secondary interactions in a QGP or a hadronic gas [16, 17]. So in this work, 3-flavour QGP emission rate [17] will be employed, and the beginning time of counting varies instead of the fixed value hydro initial time $\tau_0$. The appropriate beginning time for this counting to mimic the recent data will provide us an estimation of QGP formation $\tau_{\rm QGP}$.

The paper is organized as following: after a brief introduction of calculate approach, we present the results and the interpretation, then conclusion and discussion.

## II. APPROACH

In our previous work[13], we successfully explained the $p_{\rm t}$ spectra of direct photons from AuAu collisions measured by PHENIX with the hydrodynamical model chosen in this work, and made a detailed investigation on the competition among the main direct photon sources.

---

[*] Electronic address: liufm@iopp.ccnu.edu.cn

The collisions between hard partons and the bulk matter, make a photon source called as jet-photon conversion(JPC) whose contribution increases with system size. At the mean time, those collisions reduce the contribution of another source, fragmentation photons, or the higher order contribution of prompt photons. The increase and reduction cancel each other, so that $R_{AA} = 1$ and $v_2 = 0$ at large $p_{\rm t}$ and coincide with the measurement. Thus, we can simplify the direct photon calculation, considering neither JPC nor energy loss. Therefore, we consider only two main sources of direct photon production, prompt photons and thermal photons. The pre-equilibrium contribution to photon emission is not considered separately in the calculation. We will discuss the effect due to this approximation.

The prompt photon production is calculated as next-to-leading order cold nuclear collisions:

$$\begin{aligned} \frac{dN^{\rm P}}{dyd^2p_{\rm t}} &= T_{AB}(b)\sum_{ab}\int dx_a dx_b G_a(x_a,M^2)G_b(x_b,M^2) \\ &\times \frac{\hat{s}}{\pi}\delta(\hat{s}+\hat{t}+\hat{u})[\frac{d\sigma}{d\hat{t}}(ab\to\gamma+X) \qquad (1) \\ &+ K\sum_c \frac{d\sigma}{d\hat{t}}(ab\to cd)\int dz_c \frac{1}{z_c^2}D_{\gamma/c}(z_c,Q^2)], \end{aligned}$$

where $T_{AB}(b)$ is thickness function at given centrality, $G(x,M^2)$ is nuclear parton distribution function with nuclear shadowing effect and EMC effect, and the rest is similar to the calculation in pp collisions, *i.e.* [18].

The thermal photon calculation is based on the above-mentioned hydrodynamical description of each collision system. The solution of hydrodynamical equation will provide us the temperature $T$ and flow velocity $u_\mu$ at each space-time point of the system. Thus, the $p_{\rm t}$ spectrum of thermal photons reads

$$\frac{dN^{\rm T}}{dyd^2p_{\rm t}} = \int d^4x\Gamma(E^*,T) \qquad (2)$$

where $\Gamma(E^*,T)$ is the photon emission rate at temperature $T$, which covers emission from 3-flavour QGP and later formed hadronic gas. $E^* = p^\mu u_\mu$, $p^\mu$ is the four-momentum of a photon in the lab frame. Conventionally, $d^4x$ makes the space-time integral over the whole space and the whole emission history of the system, so the time integral begins at hydro initial time $\tau_0$. In this paper, we will try different choices. The best choice may provide an estimation to the QGP formation time $\tau_{\rm QGP}$.

The elliptic flow of thermal photons, quantified by the second harmonic coefficient

$$v_2^{\rm T}(p_{\rm t}) = \frac{\int d\phi\cos(2\phi)d^3N^{\rm T}/dy\,d^2p_{\rm t}}{\int d\phi d^3N^{\rm T}/dy\,d^2p_{\rm t}} \qquad (3)$$

comes from the Lorentz boost from the local rest frame to the lab frame. In the former frame, photons are emitted isotropically. The connection of the two frames is flow velocity $u_\mu$. So both its strength and asymmetry in the non-central collisions, are crucial for the elliptic flow of thermal photons [19].



Figure 1: (Color Online) Photons from PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, centrality 0-40%. Calculated results with four beginning times are presented as four kinds of curves. Full squares are ALICE data [1, 4]. For more details, see text. (a): Transverse momentum spectrum of direct photons. (b): Transverse momentum spectrum of thermal photons and prompt photons. (c) The elliptic flow of direct photons. (d) The elliptic flow of thermal photons.

To compare with the elliptic flow data, we should take into account of prompt photons, though they have $v_2$ zero. Thus the elliptic flow of direct photons reads

$$v_2(p_{\rm t}) = \frac{n^{\rm T}}{n^{\rm T}+n^{\rm P}}v_2^{\rm T}(p_{\rm t}), \qquad (4)$$

where $n^{\rm T}$ ($n^{\rm P}$ ) refers to $dN^{\rm T}/dyd^2p_{\rm t}$ ($dN^{\rm P}/dyd^2p_{\rm t}$).

## III. RESULTS

In Fig. 1 is shown the midrapidity photons from PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV of 0-40% centrality. Full squares are ALICE data [1, 4]. Four curves present results from four beginning times, $\tau$ =0.35 fm/c (solid line), 0.75 fm/c (dashed line), 1.15 fm/c (dotted dashed line) and 1.55 fm/c (dotted line), respectively.

In Fig. 1(a) is shown the $p_{\rm t}$ spectrum of direct photons. We can see, if we count thermal photons from $\tau_0 = 0.35$ fm/c, then the curve is higher than data. The rest three curves with delayed counting can meet the data points. This is to say, QGP is formed later than the hydro initial time and the photon emission rate during $[0.35, 0.75]$ fm/c is overestimated. The measured $p_{\rm t}$ spectrum rules out 0.35 fm/c, but can not tell which of the rest three is the best beginning time. More precise measurement like elliptic flow is needed.

In Fig. 1(b), prompt photons (black dashed line) and thermal photons (curve type same as in Fig. 1(a)) are shown separately. All the four thermal curves meet each other at low $p_{\rm t}$ region and also meet ALICE data points there. Why? Because the hydro evolution at late stage is common for the four curves, which is well controlled with bulk hadrons[10].

In Fig. 1(d) is shown the elliptic flow of thermal photons, which increase from blow to up, for the four beginning time of counting thermal photons, 0.35, 0.75, 1.15 and 1.55 fm/c, respectively. This is a clear illustration how the elliptic flow grows with time. The initial flow velocity is zero, so that the photons emitted at the first time interval have zero elliptic flow. Thus the red line decreases to zero at high $p_{\mathrm{t}}$ region.

The elliptic flow of direct photons is shown in Fig. 1(c). Because of the dominance of prompt photons, it vanishes at high $p_{\mathrm{t}}$, for all the four choices of beginning time. Also for the four curves, the increase and decrease mode and the turning point at $p_{\mathrm{t}} = 2$ GeV, agrees well with ALICE data (full squares). Among the curves, the elliptic flow can increase a few times due to the delayed counting of thermal photon emission, a much stronger effect than fluctuating the initial condition. This big increase also tells elliptic flow is more sensitive than $p_{\mathrm{t}}$ spectrum to the QGP formation time.

One may expect to meet the large elliptic flow observed by ALICE via even later beginning time. This can enlarge the elliptic flow of thermal photons, but weaken the percentage of thermal photons in eq. (4). Thus the elliptic flow of direct photons can not be increased. Besides, too late beginning may also be ruled out by the $p_{\mathrm{t}}$ spectrum.

The resulted elliptic flow of direct photons, even with delayed 3-flavour QGP photon counting, still below the data, tells us the limitation of the smoothed 3+1D ideal hydrodynamical model.

Now we turn to RHIC. In Fig. 2(a) is shown the $p_{\mathrm{t}}$ spectra of direct photons at centralities 0-20% and 20-40%, where the results of 0-20% centrality have been increased with a factor of ten for better illusion. The PHENIX results are plotted as full triangles and full squares for the two centralities, respectively. The solid lines correspond to the conventional choice, thermal photons begin to emit at $\tau_0 = 0.6$ fm/c. The simplified calculation, counting prompt photons (without energy loss) and thermal photons, does reproduce the PHENIX observed $p_{\mathrm{t}}$ spectra, as well as the consideration of four main contributions with detailed consideration of energy loss in [13]. Surprisingly, all the four kinds of curves, for four choices of beginning time such as 0.6, 1.1, 1.6 and 2.1 fm/c, are so close to each other that the measured $p_{\mathrm{t}}$ can not distinguish them.

In Fig. 2(b) and (c), is shown the elliptic flow of direct photons for centrality 0-20% and 20-40%, respectively. Similar to the case of PbPb collisions at $\sqrt{s_{NN}}$ =2.76 TeV, later counting of thermal photon emission can increase the elliptic flow. Comparing with $\tau_0 = 0.6$ fm/c, the beginning time of 2.1 fm/c makes the enhance of elliptic flow by a factor of about 3 for centrality 0-20% and 4 for centrality 20-40%! This is a much bigger effects than the initial fluctuation.

The $p_{\mathrm{t}}$-dependence of calculated $v_2$ has a similar tend as the observed data, increase first, then decrease, due to the competition between thermal photons and prompt photons. However, the turning point appears at $p_{\mathrm{t}}$=1.5 GeV/c, smaller than both PHENIX data and the value calculated for PbPb collisions at 2.76 TeV.

There are two possible reasons for the smaller turning $p_{\mathrm{t}}$. One is that the initial time of AuAu system is $\tau_0$=0.6 fm/c, later than $\tau_0 = 0.35$ fm/c in PbPb system. The big $\tau_0$ cuts the earliest growth of the flow velocity and weakens the competition of thermal photons, both in the magnitude of elliptic flow and in the slope of $p_{\mathrm{t}}$ spectrum. An early initial time of hydrodynamical expansion, can certainly improve the situation. The other reason may be the equations of state(EoS), the relation between pressure and energy density, is based on first-order transition in AuAu system, while a more realistic EoS with cross-over phase transition from lattice QCD [20], in PbPb system. A relatively long co-existence of the two phases appears in AuAu system, while no pressure to accelerate the expansion in the coexistence region.

Again, though the later thermal photon emission can increase the elliptic flow of direct photons, the best choice of beginning time, ie. 2.1 fm/c, makes the elliptic flow still below the PHENIX data points, consistent with the case of PbPb collisions at 2.76TeV.

## IV. CONCLUSION AND DISCUSSION

In conclusion, we calculated $p_{\mathrm{t}}$ spectrum and elliptic flow of direct photons from AuAu collisions at 200 GeV and PbPb collisions at 2.76 TeV, with a 3+1D event-averaged smoothing hydrodynamical description with a constrain of big hadronic database, especially the elliptic flow of hadrons. The obtained results have been compared with the recent direct photon data from RHIC and LHC.

The measured large elliptic flow of direct photons (consistent between RHIC and LHC) is still under-predicted with present hydrodynamical model. More details such as fluctuation, viscosity and earlier beginning of collective motion can improve the prediction. The earlier beginning of collective motion, *i.e.*, $\tau_0 < 0.6$ fm/c can improve the competition of thermal photons to prompt ones, as we can see from the two different initial times in the two hydrodynamical models used for AuAu collisions at RHIC energy and PbPb collisions at LHC energy.

The most important point we learned here is, a delayed thermal photon emission from 3-flavour QGP, can explain the elliptic flow much better and be allowed by measured $p_{\mathrm{t}}$ spectra of direct photons. The best beginning time to count photons from 3-flavour QGP is about 2 fm/c, later than the beginning of collective motion.

In our calculation, the pre-equilibrium contribution is not calculated separately. The elliptic flow of this contribution can not be big to solve the large elliptic flow puzzle, because of the very early emission time. In our



Figure 2: (Color Online) Direct photons from Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. PHENIX data are shown as full triangle and full squares for centrality 0-20% and 20-40%, respectively. (a): Transverse momentum spectrum of direct photons, Results of centrality 0-20% have been increased with a factor of ten for better illustration. PHENIX data from [2]. (b) and (c): Compare calculated elliptic flow $v_2$ of direct photons with measured data[3] for centrality 0-20% and 20-40%, respectively.

calculation, the beginning time of counting thermal photons is adjusted to mimic the experimental data, which cover the pre-equilibrium contribution. But the emission rate is overestimated when the equilibrium one is taken as an approximation. That's why a break of the emission between prompt photons and thermal photons appears in the calculation. As a result, this brings an error for our estimation of QGP formation time, which should be slightly later than the best beginning time we have extracted. Deeper study of the pre-equilibrium stage, with more knowledge of the matter before equilibrium, will improve our estimation of QGP formation time. And what happens before the QGP formation in the collision system, becomes an even more interesting topic. Cross checking, *i.e.* from the nuclear modification factor and elliptic flow of hadrons at high $p_t$ and heavy-flavour hadrons, should be done to test this earlier collective expansion and later formation of 3-falvour QGP in high energy nuclear collisions.

## Acknowledgments

This work is supported by the Natural Science Foundation of China under the project No. 10505010 and No. 11275081 and by Program for New Century Excellent Talents in University (NCET). The author thanks K. Werner and T. Hirano for providing the hydrodynamical evolution of two collision systems.

---

[1] M. Wilde, arXiv:1210.5958 [hep-ex].
[2] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **94**, 232301 (2005), [nucl-ex/0503003].
A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **104**, 132301 (2010), [arXiv:0804.4168 [nucl-ex]].
[3] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **109**, 122302 (2012) [arXiv:1105.4126 [nucl-ex]].
[4] D. Lohner and f. t. A. Collaboration, arXiv:1212.3995 [hep-ex].
[5] G. Basar, D. Kharzeev, D. Kharzeev and V. Skokov, Phys. Rev. Lett. **109**, 202303 (2012) [arXiv:1206.1334 [hep-ph]].
[6] S. Afanasiev *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **109**, 152302 (2012) [arXiv:1205.5759 [nucl-ex]].
[7] M. Chiu, T. K. Hemmick, V. Khachatryan, A. Leonidov, J. Liao and L. McLerran, arXiv:1202.3679 [nucl-th].
[8] R. Chatterjee, H. Holopainen, T. Renk and K. J. Eskola, arXiv:1210.3517 [hep-ph].
[9] T. Hirano, U. Heinz, D. Kharzeev, R. Lacey, and Y. Nara, Phys. Lett. B 636, 299 (2006); J. Phys. G 34, S879 (2007); Phys. Rev. C 77, 044909 (2008).
[10] K. Werner, I. .Karpenko, M. Bleicher, T. Pierog and S. Porteboeuf-Houssais, Phys. Rev. C **85**, 064907 (2012) [arXiv:1203.5704 [nucl-th]].
[11] J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757**, 102 (2005) [nucl-ex/0501009].
[12] C. Gale, arXiv:1208.2289 [hep-ph].
[13] F. -M. Liu, T. Hirano, K. Werner and Y. Zhu, Phys. Rev. C **79**, 014905 (2009) [arXiv:0807.4771 [hep-ph]].
[14] H. Holopainen, S. Rasanen and K. J. Eskola, Phys. Rev. C **84**, 064903 (2011) [arXiv:1104.5371 [hep-ph]].
[15] L. D. Landau, Izv. Akad. Nauk. SSSR **17**, 51 (1953). S. Z. Belenkij and L. D. Landau, Usp. Fiz. Nauk **56**, 309 (1955); Nuovo Cimento Suppl1. **3**, 15 (1956). S. -X. Liu, H. -J. Wang and F. -M. Liu, arXiv:1210.2833 [hep-ph].
[16] J. Kapusta, P. Lichard and D. Seibert, Phys. Rev. D **44**:2774,(1991);**47**:4171(E),(1991).
[17] P. Arnold, G. D. Moore, and L. G. Yaffe, J. High Energy Phys. **0111**, 057 (2001); J. High Energy Phys. **0112**, 9 (2001).
[18] F. -M. Liu and K. Werner, Phys. Rev. Lett. **106**, 242301 (2011) [arXiv:1102.1052 [hep-ph]].
[19] F. -M. Liu, T. Hirano, K. Werner and Y. Zhu, Phys. Rev. C **80**, 034905 (2009) [arXiv:0902.1303 [hep-ph]].
[20] S. Borsanyi, G. Endrodi, Z. Fodor, A. Jakovac, S. D. Katz, S. Krieg, C. Ratti and K. K. Szabo, JHEP

**1011**, 077 (2010) [arXiv:1007.2580 [hep-lat]].